

your name _____

Physics 321 Midterm #2 - Wednesday, Nov. 9, 2022

Some integrals:

$$\begin{aligned}\int \frac{dx}{1+x^2} &= \tan^{-1}(x), \\ \int \frac{dx}{1-x^2} &= \tanh^{-1}(x), \\ \int \frac{dx}{\sqrt{1-x^2}} &= \sin^{-1}(x), \\ \int \frac{dx}{\sqrt{1+x^2}} &= \sinh^{-1}(x), \\ \int dx \tan(x+\phi) &= -\ln(\cos(x+\phi)) \\ \int dx \tanh(x+a) &= \ln(\cosh(x+a)).\end{aligned}$$

$$\begin{aligned}F(t) &= \frac{f_0}{2} + \sum_n f_n \cos(n\omega t) + g_n \sin(n\omega t), \\ f_n &= \frac{2}{\tau} \int_{-\tau/2}^{\tau/2} dt F(t) \cos(2n\pi t/\tau), \\ g_n &= \frac{2}{\tau} \int_{-\tau/2}^{\tau/2} dt F(t) \sin(2n\pi t/\tau).\end{aligned}$$

For the equation

$$\ddot{x} + 2\beta\dot{x} + \omega_0^2 x = 0,$$

The solutions are:

$$\begin{aligned}x &= Ae^{i\omega t}, \\ \omega &= i\beta \pm \sqrt{\omega_0^2 - \beta^2}, \\ x(t) &= Ae^{-\beta t} e^{\pm i\sqrt{\omega_0^2 - \beta^2} t}, \text{ underdamped,} \\ &= Ae^{-\beta t \pm \sqrt{\beta^2 - \omega_0^2} t}, \text{ overdamped.}\end{aligned}$$

Write your name on EVERY page!

80 points possible

your name _____

1. (15 pts) A mass m is connected to a spring with spring constant $k = m\omega_0^2$ and has a very small damping. The mass also feels a periodic force

$$F(t) = F_0 \sin(\omega t),$$
$$\omega \neq \omega_0.$$

Assuming the force has been applied for a very very long time, find the displacement $x(t)$ for $t > 0$.

your name _____

Extra Space for No.1

your name _____

2. A mass m is connected to a spring with spring constant $k = m\omega_0^2$ and has an very small damping constant. It is subject to a periodic force,

$$F(t) = \begin{cases} -F_0, & -\tau/2 < t < 0, \\ F_0, & 0 < t < \tau/2 \end{cases},$$
$$F(t + \tau) = F(t).$$

- (a) (15 pts) Expressing the force as a series

$$F(t) = \frac{f_0}{2} + \sum_n f_n \cos(n\omega t) + g_n \sin(n\omega t),$$
$$\omega = 2\pi/\tau,$$

what are the coefficients f_n and g_n ? Note that $\omega \neq \omega_0$.

- (b) (10 pts) Assuming the force has been active for a very very long time, express the displacement $x(t)$ in terms of the coefficients (as a sum) .

your name _____

Extra Space for No.2

your name _____

3. Consider a particle of mass m and energy E which experiences an attractive spherically symmetric potential,

$$V(r) = -\frac{V_0}{r^2}.$$

If the particle reaches $r = 0$ it will annihilate.

- (a) (5 pts) Sketch the effective (real plus centrifugal) potential for a particle with angular momentum L . Show potentials for a few values of L to demonstrate how V_{eff} depends on L .
- (b) (15 pts) Calculate the cross section for annihilation.

your name _____

Extra Space for No.3

your name _____

4. (20 pts, no points awarded if answer has wrong dimension) You are an experimentalist planning a scattering experiment using a gold target and a much lighter projectile. You go online and find the mass density of gold, ρ_{Au} , and the mass of a gold atom, m_{Au} . Somebody sends you a target, but they didn't tell you the dimensions of the thin foil. You measure the width and length, W and L , then measure the mass of the foil, M , but you can't measure the thickness. The accelerator provides a beam, with N_{beam} particles incident on the foil over the duration of the experiment. One of your detectors is positioned at a scattering angle $\theta_s = 45^\circ$. The detector has a cross-sectional area a and is placed a distance R from the center of the target. Over the six days you record N_{counts} in the detector. What is $d\sigma/d\Omega$ for $\theta_s = 45^\circ$?

your name _____

Extra Space for No.4