

your name(s) \_\_\_\_\_

---

Physics 852 Exercise #4b - Friday, Feb. 4th

The radial hydrogen-atom wave functions are

$$\psi_{n,\ell}(r, \theta, \phi) = \left\{ \left( \frac{2}{na_0} \right)^3 \frac{(n - \ell - 1)!}{2n[(n + \ell)!]^3} \right\}^{1/2} e^{-r/(na_0)} \left( \frac{2r}{na_0} \right)^\ell L_{n+\ell}^{2\ell+1} \left( \frac{2r}{na_0} \right) Y_{\ell,m}(\theta, \phi).$$

Here,  $L_n^k(x)$  are Laguerre polynomials.

Consider the operator

$$A \equiv e^{-R^2/R_0^2} X^2,$$

where  $R_0$  is some arbitrary constant. The operators  $X, Y, Z$  are the position operators and  $R^2 = X^2 + Y^2 + Z^2$ .

1. Write the operator  $X^2$  in terms of a sum over irreducible tensor operators,  $T_q^k$ , where you define the operators. (You can find this in the lecture notes or peek at the FYI below).
2. You need to calculate the matrix elements

$$\langle n, \ell, m | A | 0 \rangle.$$

For which values of  $n, \ell, m$  might the matrix element be non-zero? Use the orthogonality properties of spherical harmonics.

3. Repeat (2) but replace  $A$  with the operator

$$B \equiv e^{-R^2/R_0^2} P_x^2,$$

where  $P_x, P_y, P_z$  are the momentum operators. You may wish to use the expression for  $P_x^2$  in spherical coordinates given in the FYI.

4. After completing 1-3, go home and think about how you would go about answering the same questions but with the ket being  $|n', \ell', m'\rangle$ . Don't do it!

FYI: Some spherical harmonics are:

$$\begin{aligned}
 Y_{0,0} &= \frac{1}{\sqrt{4\pi}}, \\
 Y_{1,0} &= \sqrt{\frac{3}{4\pi}} \cos \theta, \\
 Y_{1,\pm 1} &= \mp \sqrt{\frac{3}{8\pi}} \sin \theta e^{\pm i\phi}, \\
 Y_{2,0} &= \sqrt{\frac{5}{16\pi}} (3 \cos^2 \theta - 1), \\
 Y_{2,\pm 1} &= \mp \sqrt{\frac{15}{8\pi}} \sin \theta \cos \theta e^{\pm i\phi}, \\
 Y_{2,\pm 2} &= \sqrt{\frac{15}{32\pi}} \sin^2 \theta e^{\pm 2i\phi}, \\
 Y_{\ell-m}(\theta, \phi) &= (-1)^m Y_{\ell m}^*(\theta, \phi).
 \end{aligned}$$

Using the following definition of some irreducible tensor operators,

$$\begin{aligned}
 T_0^0 &= 1, \\
 T_1^1 &= -\frac{1}{\sqrt{2}}(x + iy), \\
 T_0^1 &= z, \\
 T_{-1}^1 &= \frac{1}{\sqrt{2}}(x - iy), \\
 T_2^2 &= \sqrt{\frac{3}{8}}(x^2 + 2ixy - y^2), \\
 T_1^2 &= -\frac{\sqrt{3}}{2}z(x + iy), \\
 T_0^2 &= \frac{1}{2}(3z^2 - r^2), \\
 T_{-1}^2 &= \frac{\sqrt{3}}{2}z(x - iy), \\
 T_{-2}^2 &= \sqrt{\frac{3}{8}}(x^2 - 2ixy - y^2),
 \end{aligned}$$

one can express various powers of  $x$ ,  $y$  and  $z$ .

$$\begin{aligned}
1 &= T_0^0, \\
x &= \frac{1}{\sqrt{2}}(T_{-1}^1 - T_1^1), \\
y &= \frac{i}{\sqrt{2}}(T_{-1}^1 + T_1^1), \\
z &= T_0^1, \\
x^2 &= \frac{1}{2}\sqrt{\frac{2}{3}}(T_2^2 + T_{-2}^2) - \frac{1}{3}T_0^2 + \frac{1}{3}T_0^0 r^2, \\
y^2 &= -\frac{1}{2}\sqrt{\frac{2}{3}}(T_2^2 + T_{-2}^2) - \frac{1}{3}T_0^2 + \frac{1}{3}T_0^0 r^2, \\
z^2 &= \frac{2}{3}T_0^2 + \frac{1}{3}T_0^0 r^2, \\
xy &= i\frac{1}{\sqrt{6}}(T_{-2}^2 - T_2^2), \\
xz &= \frac{1}{\sqrt{3}}(T_{-1}^2 - T_1^2), \\
yz &= \frac{i}{\sqrt{3}}(T_{-1}^2 + T_1^2).
\end{aligned}$$

Some algebra,

$$\begin{aligned}
\partial_x f(r, \theta, \phi) &= \frac{\partial r}{\partial x} \partial_r f + \frac{\partial \theta}{\partial x} \partial_\theta f + \frac{\partial \phi}{\partial x} \partial_\phi f \\
&= \frac{\sin \theta \cos \phi}{r} \partial_r f + \frac{\cos \theta \cos \phi}{r} \partial_\theta f - \frac{\sin \phi}{r \sin \theta} \partial_\phi f
\end{aligned}$$

If  $f(r, \theta, \phi)$  depends only on  $r$ ,

$$\begin{aligned}
\partial_x^2 f(r) &= \left( \frac{\sin \theta \cos \phi}{r} \partial_r + \frac{\cos \theta \cos \phi}{r} \partial_\theta - \frac{\sin \phi}{r \sin \theta} \partial_\phi \right) \frac{\sin \theta \cos \phi}{r} \partial_r f \\
&= \frac{\sin^2 \theta \cos^2 \phi}{r^2} \partial_r^2 f + (\cos^2 \theta \cos^2 \phi + \sin^2 \phi) \frac{1}{r} \partial_r f \\
&= \frac{\sin^2 \theta \cos^2 \phi}{r^2} \partial_r^2 f + (1 - \sin^2 \theta \cos^2 \phi) \frac{1}{r} \partial_r f
\end{aligned}$$