



Integrative Studies in Physical Science ISP209

Fall Semester 2005
December 1 Lecture
Section 001

Professor Stan Schriber

The Mystery of the Physical World



- **ISP 209**
 - *Accelerator Physics today!*
 - *No Quiz*
- **Homework?**
 - *Two problems*
- **DOCS:**
 - *Lecture VGs one day early*
- **Questions?**




What is an Accelerator




- A device that speeds particles to high velocities.
 - Types:
 - DC
 - RF
 - DC Examples
 - X-ray Tubes
 - Ion Sources
 - Van de Graff
 - Tandem Accelerator
 - RF Examples
 - Linear machines
 - Circular Machines

DC Devices are Limited


- Voltage standoff is biggest problem - but very useful devices
 - Limits velocity attainable
- Sterilization
- X-rays
- Food irradiation
- Implantation
- Research




DC Example - Tandem Accelerator



Oxygen Beam Planter




750 keV Cockcroft-Walton




Sits in a Faraday Cage; components with rounded edges to minimize arcs

Radio Frequency (RF) Wave



RF Field in Cavity as a Function of Time




Drift Tube Linac



Drift Tube Linac

- The reason for drift tubes with an oscillating rf system




- Fields when in the wrong direction for acceleration are shielded by the drift tubes.
- Particles get accelerated at each gap between drift tubes.

Cavity Examples



A voltage generator induces an electric field inside the rf cavity. Its voltage oscillates with a radio frequency of 1.3 Gigahertz or 1.3 billion times per second.

The electrons always feel a force in the forward direction.




An electron source injects particles into the cavity in phase with the variable voltage.


The electrons never feel a force in the backward direction.



Coupled Cavity



Coupled Cavity



Relativistic Effects

$$\beta = v/c$$

$$\gamma = \frac{1}{\sqrt{1-\beta^2}}$$

$$E_{KE} = m_0 c^2 (\gamma - 1) = mc^2$$

$$m_0 c^2 \approx 0.511 \text{ MeV} \quad \text{for an electron}$$

$$m_0 c^2 \approx 938 \text{ MeV} \quad \text{for a proton}$$

$$\text{Length contraction: } L = L_0 / \gamma$$

$$\text{Time dilation: } \Delta t = \gamma \Delta t_0$$




Fig. 3. Graph of the variation of the velocity and mass of electrons and protons as a function of their kinetic energy.

Why Colliding Beams



- Colliding beams are much more effective in obtaining a high energy for the collision in the center of mass frame as compared to colliding on a fixed target.

- **Colliding beam effective energy**

$$E \approx 2\sqrt{\gamma_1\gamma_2}m_0c^2$$

- 10GeV proton on 10GeV proton is ~20 GeV effective

- **Fixed target effective energy**

$$E \approx \sqrt{2\gamma_1}m_0c^2$$


- Need ~200GeV proton on proton target to get 20GeV effective

WOW! Really hard in comparison

Circular Machines


- Match centripetal force with magnetic force on the ions to keep particles in a circular orbit.

$$\frac{mv^2}{r} = qvB$$




Particle with mass m (kg) and positive charge q (C) moving at velocity v (m/s) in a magnetic field pointing into the page of intensity B (T) follows a circular orbit of radius r (m).

Types of RF Accelerators




Linacs



Circular

- *Betatron*
- *Microtron*
- *Cyclotron*
- *Synchrotron*
- *Storage Ring*
- *Accumulator Ring*

High Power Linac Basics





Many important components not shown: rf, cooling, controls, diagnostics, chopper, support, magnets, power supplies, kickers, etc.

Maintain focusing periodicity throughout (soft dough and F0D0)


Ion Linac Necessity - RFQ

- Radio Frequency Quadrupole (RFQ) **1979**
 - Matches, Bunches, Gently & Adiabatically, Accelerates



Focusing in a DTL

F – Focus (Blue)
D – Defocus (Red)
Accelerate (Green)




e⁻ and p Linacs



- Cancer Therapy (1000's)
- Structural Investigation
- Oil well logging
- Isotope Production
- PET systems
- MRI
- Sterilization

Schematic diagram of a typical medical accelerator used in cancer radiotherapy.



- Proton/neutron therapy



Linac Injectors





LiNSTAR™ 

© AccSys Technology, Inc.



Loma Linda Synchrotron




■ Proton therapy



U of Tsukuba 250 MeV



Superconducting Cyclotron for Neutron Therapy (Detroit - MSU)



Superconducting Cyclotrons

- Movie from Nova program "The Nucleus Factory"



K500



K1200 +
K500

Stanford Linear Accelerator Center

SLAC (1)



- ON THE CHILLY SPRING evening of April 10, 1956, a number of boxy Fords and Chevrolets began pulling up in front of the rambling, ranch-style Los Altos Hills home of a young Stanford physics professor named Wolfgang K. H. Panofsky. About 20 earnest-looking young and middle-aged men with short-cropped hair got out and walked into the house. Neighbors thought nothing of this convergence-- Panofsky had been hosting regular Monday night bull sessions with students for several years. But this was not a Monday night, and it certainly was no ordinary bull session.
- Gathered in Panofsky's living room that night were the top professors in Stanford's electrical engineering, microwave and high-energy physics laboratories. For these normally cautious men of science, the concept under discussion was breathtaking. "All other physical sciences, and probably all life sciences, must ultimately rest on the findings of elementary particle physics," Panofsky, known since childhood as "Pief," would later write. "We cannot afford to be ignorant of the most fundamental type of structure on which everything else depends." On this April night, these men were setting out on a quest to find that fundamental structure--the basic building blocks of the universe.
- Encouraged by early experiments on subnuclear matter obtained using the University's 220-foot long Mark III electron accelerator, Panofsky and the others had begun dreaming about a massive scale-up. Their audacious vision: a machine that would generate 50 times the power of the Mark III and extend in a straight line over two full miles. As physics professor and Nobel laureate Felix Bloch, an initial skeptic, later told Panofsky, "Pief, if you must build a monster, build a good monster."

SLAC (2)



SLAC (3)



1962

SLAC Ground Breaking

1966

Linac Begins Operation

1968

Quarks Discovered in Nucleon

Nobel Prize 1990

1974-76


Charm Discovered at SPEAR

Nobel Prize 1976

1976-78

Tau Lepton Discovered at SPEAR

Wolf Prize 1983; Nobel Prize 1995



1980-82

B Meson Lifetime Measured at PEP

1989-90

Limit of Three Quark Generations
Measured at SLC

1991

Operation with Polarized Zs at SLC/SLD

1994

Construction Begins on (PEP-II)
B-Factory

Alternating Gradient Synchrotron



AGS at BNL



- In the early 1950's, scientists knew that achieving the higher energies needed for future research was going to be a difficult problem. Calculations showed that, using existing technology, building a proton accelerator ten times more powerful than the 3.3-billion electron volt (GeV) Cosmotron would require 100 times as much steel. Such a machine would weigh an astronomical 200,000 tons. Brookhaven physicists Ernst Courant, M. Stanley Livingston, and Hartland Snyder overcame this barrier by co-inventing the alternating gradient or strong-focusing principle of propelling protons.




AG Big Break Through

- In the Cosmotron, all the magnets were C-shaped, with the open side and the magnetic field, facing outward. The breakthrough occurred by alternating the orientation of these magnets, so some of their field gradients faced outward and some inward. Brookhaven physicists found that the net effect of alternating the field gradient was that both the vertical and horizontal focusing of protons could be made strong at the same time, allowing tight control of proton paths in the machine (right). This increased beam intensity while reducing the overall construction cost of a more powerful accelerator.




Magnets


Dipole Extraction/Injection




Quadrupole



Alternating Gradient Concept



Octupole




Sextupole


Magnet Issues

- Synchrotron radiation
- Beam pipe and vacuum
- Superconducting coils
- Radiation environment
- Enormous forces from magnetic fields
- Materials and alignment


RHIC (1)



The Alternating Gradient Synchrotron complex



RHIC (2)



Siberian Snake



- FNAL - Tevatron *Coupled Cavity Linac*




Drift Tube Linac

Dipole Construction



Tevatron (2)



CERN (1)



CERN in numbers


- Financed by 20 European countries
 - Special contributions also from other countries:
 - USA, Canada, China, Japan, Russia, etc.
- 1000 CHF (650 M€) budget to cover operation + new accelerators
- 2,200 staff (and diminishing)
- 6,000 users (researchers) from all over the world
 - broad visitor and fellowship program

May 2004

Svenja Jarp



Collisions at the Large Hadron Collider




Introduction to CERN

David Barney, CERN



LHC Detectors

General-purpose
Higgs
SUSY
??




Introduction to CERN


David Barney, CERN



Accelerators and detectors in underground tunnels and caverns



CERN Beam Gymnastics (2)




CERN (3)






What is CERN?

- Physicists smash particles into each other to:
 - identify their **components**
 - **create** new particles
 - reveal the nature of the **interactions** between them
 - create an environment similar to the one present at the origin of our Universe
- **What for?** To answer fundamental questions like:
how did the Universe begin? What is the origin of mass? What is the nature of antimatter?




CERN Site






LHC DIPOLE : STANDARD CROSS-SECTION



CERN AC/DI/MMA - HE107 - 30 04 1999



1st hardware for LHC from USA (5)



CERN (6)



Map of CERN sites and LHC access points



RIA Facility



- Rare Isotope Accelerator

