



Today

- Announcements:
 - The average on the first exam was 31/40
 - Exam extra credit is due by 8:00 am Wednesday February 13th.
 - The second homework extra credit is due by Wednesday February 13th.
- F=ma
- Electric Force
- Work, Energy and Power



Newton's Second Law of Force

F=ma


An equation is worth greater than half the words of a picture.

- Force is equal to mass times acceleration.
- For a given force, the amount of acceleration is inversely proportional to the mass.
- Force causes acceleration.
- If you observe acceleration, there must be a force acting.



A new Force!

- Charge is a property of matter. It is measured in Coulombs C.
- Like charges repel, unlike charges attract.
- Coulomb's Law of Electric Force



$$F = \frac{kQ_1Q_2}{r_{12}^2} \quad k = 8.99E9 \text{ N} \cdot \text{m}^2 / \text{C}^2$$



Why?

- Coulomb's law looks like Newton's Law of gravity. Why?
- Why does charge come in two types and mass only came in one type?
- Why do we always get r²? I hate squares.
- Why is k = 8.99E9 Nm²/C² so much bigger than G = Nm²/kg²?

Two possible answers:

- (1) I can't tell you until you are older. (2) I don't know.



Energy

- Energy is the ability to do work
- Energy comes in two forms
 - Kinetic (KE) – energy of motion
 - Potential (PE) – energy of position
- There are many variants on these type main types, e.g. chemical, nuclear, thermal, ...



Energy and Power

- **Energy is the ability to do work:** Work = force x distance = $F d$
- Energy comes in two forms
 - Kinetic (KE) – energy of motion
 - Potential (PE) – energy of position

Gravitational GPE = $m (gh)$; $g = 9.81 \text{ m/s}^2$ on Earth, h height
- **Power** (measured in $W = \text{J/s}$) is the rate of change (or use) of energy

$$KE = \frac{1}{2}mv^2$$

m - mass
 v - velocity



Some Example Problems

Examples:


- A mass of 1.0 kg is raised 1.0 m. How much work was done?
 $W = \Delta GPE = mg\Delta h = 1.0 \text{ kg} \times 9.81 \text{ m/s}^2 \times 1.0 \text{ m} = 9.81 \text{ J}$
- A 90.0 kg ISP209 professor walks up two flights of stairs. How much did his/her potential energy increase? DATA 1 flight of stairs = 3.00 m
 $\Delta GPE = 90.0 \text{ kg} \times 9.81 \text{ m/s}^2 \times 2 \text{ flights} \times (3 \text{ m/flight}) = 5.29 \text{ kJ}$



Conservation of Energy

In nature certain quantities are “conserved”. Energy is one of these quantities. Charge is another.

Example: Ball on a hill



A 1.00 kg ball is rolled toward a hill with an initial speed of 5.00 m/s. If the ball rolls without friction, how high, h , will the ball go?

$$KE = \frac{1}{2}mv^2 \quad PE = mgh; \quad g = 9.80 \frac{m}{s^2}$$

$$\frac{1}{2}mv^2 = mgh \rightarrow h = \frac{v^2}{2g} = \frac{(5 \text{ m/s})^2}{2 \cdot 9.80 \frac{m}{s^2}} = 1.28 \text{ m}$$



Work

- Work = Force x distance
- Work is a scalar and is measured in Joules, J
- Bill pushes on a wall with 10 N for 33 s. If the wall does not move, how much work is done on the wall?
- Work = 10.0 N x 0.0 m = 0.0 N
- How does that make sense? Work has a strict definition. If the kinetic or potential energy of the wall did not change, no work was done on the wall.
- Work changes energy from one form to another.



Power

- Power is the rate of change of energy
- Power = (change in energy)/(change in time)
- Power is a scalar and is measured in watts.
- Light bulbs are measured in watts
- Sun (a big light bulb) - 3.827×10^{26} W



Information

- Horsepower $746 \text{ W} = 1 \text{ horsepower}$
*In fourteen hundred and ninety-two
Columbus sailed the ocean blue.
And if you divide by two
You get watts in a horsepower too*
- Food energy is measured in kcal
 - 1 food cal = 4.184 J
 - 1 Calorie = 1 kcal (what we call calories are actually kilocalories)



Example Problem

How many kcal are burned by doing 1500 J of work?

DATA: The human body is 10% efficient in converting food energy to work.

$$\text{cal} = \text{energy} \cdot \frac{1 \text{ cal}}{4.184 \text{ J}} \cdot \left(\frac{1}{\text{efficiency}} \right)$$

$$1500 \text{ J} \cdot \frac{1 \text{ cal}}{4.184 \text{ J}} \cdot \left(\frac{1}{0.1} \right) = 3590 \text{ cal} = 3.59 \text{ kcal}$$